

Suggested Schedule for Adult Vaccinations

COVID-19 vaccines are recommended for everyone ages 6 months and older

Vaccine	19-26 years	27-49 years	50-64 years	≥65 years
Influenza	1 dose annually			
Respiratory Syncytial Virus (RSV)	Seasonal administration during pregnancy			1 dose
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management			
	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication			
Varicella	2 doses (if born in 1980 or later)		2 doses	
Zoster recombinant (RZV)	2 doses for immunocompromising conditions		2 doses	
Human papillomavirus (HPV)	2 or 3 doses depending on age	27 through 45 years		
Pneumococcal				Connect with your primary care team
Hepatitis A	2, 3, or 4 doses depending on vaccine			
Hepatitis B	2, 3, or 4 doses depending on vaccine or condition			
Meningococcal A, C, W, Y	1 or 2 doses depending on indication, contact your primary care team for booster recommendations			
Meningococcal B	2 or 3 doses depending on vaccine and indication, contact your primary care team for booster recommendations			
Haemophilus influenzae type b (Hib)	1 or 3 doses depending on indication			

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection.

Recommended vaccination for adults with an additional risk factor or another indication.

Recommended vaccination based on shared clinical decision-making.